

Tunneling into the vortex state of NbSe₂ with van der Waals junctions

T. Dvir,¹ M. Aprili,² C. H. L. Quay,² and H. Steinberg¹

12

FIG. 1. Spectral response of NbSe₂

FIG. 2. Sub-gap spectra at perpendicular magnetic field (a.) $G_0=G_N$ of Junctions 1,3,4,6,7 as a function of H_{\perp} normalized to the critical field $H_{c2}(T = 0)$. Dashed lines shows are a linear curve $G_0=G_n = 4H_{\perp}/H_{c2}$, that agrees with the low field dependence and a square root curve $G_0=G_n =$

[13] Eskildsen, M. R. *et al.* Vortex imaging in magnesium diboride with H₂c. *Physical Review B* 68, 63–4 (2003).

[14] Tinkham, M. *Introduction to superconductivity* (Courier Corporation, 2004).

[15] Nakai, N., Ichioka, M. & Machida, K. Field Dependence of Electronic Specific Heat in Two-Band Superconductors. *Journal of the Physical Society of Japan* 71, 23–26 (2002).

[16] Eskildsen, M. R. *et al.* Vortex Imaging in the σ Band of Magnesium Diboride. *Physical Review Letters* 89, 179–4 (2002).

[17] Kiss, T. *et al.* Charge-order-maximized momentum-dependent σ band in MgB₂. *Physical Review Letters* 144, 095501 (2020).

